

The b-chromatic index of a graph

Marko Jakovac*

University of Maribor

Faculty of Natural Sciences and Mathematics

Koroška cesta 160, 2000 Maribor, Slovenia

marko.jakovac@uni-mb.si

Iztok Peterin

University of Maribor

Faculty of Electrical Engineering and Computer Science

Smetanova ulica 17, 2000 Maribor, Slovenia

iztok.peterin@uni-mb.si

May 25, 2013

Abstract

The b-chromatic index $\varphi'(G)$ of a graph G is the largest integer k such that G admits a proper k -edge coloring in which every color class contains at least one edge incident to some edge in all the other color classes. The b-chromatic index of trees is determined and equals either to a natural upper bound $m'(T)$ or one less, where $m'(T)$ is connected with the number of edges of high degree. Some conditions are given for which graphs have the b-chromatic index strictly less than $m'(G)$, and for which conditions it is exactly $m'(G)$. In the last part of the paper regular graphs are considered. It is proved that with four exceptions, the b-chromatic number of cubic graphs is 5. The exceptions are K_4 , $K_{3,3}$, the prism over K_3 , and the cube Q_3 .

Key words: b-chromatic index, regular graphs, trees;

AMS subject classification (2010): 05C15, 05C76

1 Introduction and preliminaries

A *b-vertex coloring* of a graph G is a proper vertex coloring of G such that each color class contains a vertex that has at least one vertex in every other color class in its neighborhood. The *b-chromatic number* $\varphi(G)$ of a graph G is the largest integer k

*Corresponding author

for which G has a b-vertex coloring with $\varphi(G)$ colors. This concept was introduced in [15] by Irving and Manlove by the certain partial ordering on all proper colorings in contrast to chromatic number $\chi(G)$. Namely, $\chi(G)$ is the minimum of colors used among all minimal elements of this partial ordering, while $\varphi(G)$ is the maximum of colors used among all minimal elements of the same partial ordering.

Since then the b-chromatic number has drawn quite some attention among the scientific community. Already Irving and Manlove [15] have shown, that computing $\varphi(G)$ is an *NP*-complete problem in general. Hence an approximation approach described in [7] seems natural. The b-chromatic number of some special graph classes has been studied in [9, 8, 10, 3]. The bounds for the b-chromatic number have been studied in [19] in general and for some graph classes in [6, 21, 1, 2]. The b-chromatic number has been considered with respect to subgraphs in [12, 18], while the b-chromatic number under graph operations was considered in [20] for the Cartesian product and in [17] for the other three standard products. In [4] an interesting concept of *b-continuous graphs* was introduced as graphs for which there exists a t -b-vertex coloring for every integer t between $\chi(G)$ and $\varphi(G)$.

Intuitively, we need to have enough vertices of high enough degree, at least one in each color class. Let v_1, \dots, v_n be such a sequence of vertices, that $d(v_1) \geq \dots \geq d(v_n)$. Then $m(G) = \max\{i : d(v_i) \geq i - 1\}$ is an upper bound for $\varphi(G)$. From this point of view, d -regular graphs are of special interest, since $m(G) = d + 1$ for a d -regular graph G and every vertex is a candidate to have each color class in its neighborhood. Indeed, in [22] it was shown that if a d -regular graph G has at least d^4 vertices, then the equality $\varphi(G) = d + 1$ holds. This bound was later improved to $2d^3$ in [5]. In particular it was shown in [16], that there are only four exceptions among cubic graphs with $\varphi(G) < 4$, one of them being the Petersen graph.

We introduce in this work an edge version of the b-vertex coloring and the b-chromatic number, namely the b-edge coloring and the b-chromatic index, respectively. It is a natural approach to study vertex concepts on edges and vice versa. The classical example is the chromatic number and its edge version the chromatic index. But we can also find more recent dates. For instance in [14] the edge Wiener index was defined in four different ways. One of these approaches, via the line graph, we also use here.

A *b-edge coloring* of a graph G is a proper edge coloring of G such that each color class contains an edge that has at least one incident edge in every other color class and the *b-chromatic index of a graph G* is the largest integer $\varphi'(G)$ for which G has a b-edge coloring with $\varphi'(G)$ colors. An edge e of color i that has all other colors on its incident edges is called *color i dominating edge* or we say that color i is *realized* on e . There is no evidence for any publications regarding the b-chromatic index, but we managed to find the manuscript [23] in which authors show that this problem is NP-complete.

In the rest of this section we recall some standard notation that will be used later. In the second section we first describe some bounds, compute $\varphi'(T)$ for every

tree T , and show that general bounds with respect to $\Delta(G)$ cannot be improved. In the third section we show that for many regular graphs $\varphi'(G)$ attains the trivial upper bound. Next section is devoted to graphs where $\varphi'(G)$ is strictly less than the trivial upper bound and in the last section we describe the complete list of cubic graphs for which the trivial upper bound is not achieved.

Let G be a graph. The *line graph* $\mathcal{L}(G)$ of a graph G is the graph with $V(\mathcal{L}(G)) = E(G)$, and two edges of G are adjacent in $\mathcal{L}(G)$ if they share a common vertex. Clearly $\varphi'(G) = \varphi(\mathcal{L}(G))$. The number of vertices incident with the vertex v is called the *degree* of v and is denoted by $d(v)$. If all vertices have the same degree d , we say that G is a d -regular graph.

The *Cartesian product* $G \square H$ of graphs G and H has the vertex set $V(G) \times V(H)$. Two vertices (g, h) and (g', h') are adjacent if they are adjacent in one coordinate and equal in the other, i.e. $g = g'$ and $hh' \in E(H)$ or $gg' \in E(G)$ and $h = h'$. The Cartesian product is associative (see [11]) and hence we can write more factors without brackets: $G_1 \square \dots \square G_k$. If every factor G_i , $i \in \{1, \dots, k\}$, is isomorphic to a complete graph, then we call such a Cartesian product a *Hamming graph*. For more about Cartesian product graphs in general or Hamming graphs in particular see the books [11, 13].

2 Bounds, trees, and realizability

Every proper edge coloring of a graph G with $\chi'(G)$ colors is also a b-edge coloring. If not, then every edge of a color class with no realizable edge can be recolored by some other color. This yields a proper edge coloring with less than $\chi'(G)$ colors, which is a contradiction. Hence $\chi'(G)$ is the trivial lower bound of $\varphi'(G)$.

Let $e = uv$ be an edge of G . Denote with $N_u(e)$ the set of edges in G different from e that share u with e , and $N_v(e)$ the set of edges in G different from e that share v with e . By $N(e)$ we denote the *neighborhood of an edge* e , which is $N(e) = N_u(e) \cup N_v(e)$. The degree of an edge e is denoted with $d(e)$ and is equal to $|N(e)|$. A (realizable) edge e can have at most $2\Delta(G) - 2$ colors in its neighborhood ($\Delta(G) - 1$ in every end vertex). Together with the color of e , this gives a trivial upper bound for $\varphi'(G)$, namely $2\Delta(G) - 1$. The trivial upper bound is meaningful only if there exist enough ($\geq 2\Delta(G) - 1$) edges in G of degree $2\Delta(G) - 2$, such that every color is realized. Otherwise we can lower the trivial upper bound as follows. Let e_1, \dots, e_m be edges of G and $d(e_1) \geq \dots \geq d(e_m)$ the degree sequence of these edges. Then $m'(G) = \max\{i : d(e_i) \geq i - 1\}$ is an improved upper bound for $\varphi'(G)$. Hence

$$\Delta(G) \leq \chi'(G) \leq \varphi'(G) \leq m'(G) \leq 2\Delta(G) - 1.$$

Note that for regular graphs $m'(G) = 2\Delta(G) - 1$ and for stars $K_{1,\ell}$ we have $m'(G) = \Delta(G)$. Every realizable edge of a b-edge coloring must have degree at least $m'(G) - 1$. Hence we call an edge e with $d(e) \geq m'(G) - 1$ a *dense* edge.

The distance between edges $e, f \in E(G)$ is defined as the number of edges on a shortest path between e and f (excluding e and f).

We now present the exact value for the b-chromatic index of trees. We use a similar approach than the one used by Irving and Manlove in [15] for the b-chromatic number of trees.

Definition 2.1 *A tree T is called pivoted if it has exactly $m'(T)$ dense edges and an edge e such that:*

1. *Edge e is not dense.*
2. *There exist two dense edges that are incident with edge e and are not incident with each other.*
3. *Each dense edge is incident to e or to a dense edge incident to e .*
4. *Any dense edge incident to e and to another dense edge which is not incident to e has degree $m'(T) - 1$.*

We call e the pivot of T (for an example see Figure 1).

Clearly, the pivot is unique if it exists.

Theorem 2.2 *If T is a pivoted tree then $\varphi'(T) = m'(T) - 1$.*

Proof. Let e be the pivot of the tree T and $E(T) = \{e_1, \dots, e_{|E(T)|}\}$ its edges. Let $E'(T) = \{e_1, \dots, e_{m'(T)}\}$, $m'(T) \leq |E(T)|$, be dense edges of T . In addition, let e_1, \dots, e_p , $p \leq m'(T)$, be dense edges incident with e and e_1, \dots, e_q , $q \leq p$, dense edges incident with e that have at least one dense edge not incident to e as a neighbor. Clearly $p \geq 2$ by Property 2 of Definition 2.1 and $q \geq 1$, since e is not a dense edge.

Let us first show that $\varphi'(T) < m'(T)$. Suppose that there exists a b-edge coloring c of T with $m'(T)$ colors. Without loss of generality assume that $c(e_i) = i$, $i \in \{1, \dots, m'(T)\}$. Since $d(e_j) = m'(T) - 1$, $j \in \{1, \dots, q\}$, edges e_1, \dots, e_q are incident to exactly one edge of any other color. Moreover, every dense edge $e_{p+1}, \dots, e_{m'(T)}$ is incident with exactly one edge from e_1, \dots, e_q . Now e cannot receive color $j \in \{1, \dots, p\}$. Also it cannot receive color $j \in \{p+1, \dots, m'(T)\}$, or else some e_k , $k \in \{1, \dots, q\}$, is incident to two edges of that color and is not realizable. Hence there is no available color for e .

Next we prove that $\varphi'(T) = m'(T) - 1$ by constructing a proper b-edge coloring c of T with $m'(T) - 1$ colors. By Property 2 of Definition 2.1 there exist two edges e_1 and e_i , $2 \leq i \leq p$, which are incident to e and are not incident with each other. Also for some $r \in \{q+1, \dots, m'(T)\}$ there exists a dense edge e_r which is incident to e_1 but not to e . Set $c(e_j) = j - 1$, $j \in \{2, \dots, m'(T)\}$, $c(e) = r - 1$, and $c(e_1) = i - 1$. With this all dense edges are colored together with e .

Next we consider uncolored edges that are incident to some dense edge e_i , $i \in \{1, \dots, m'(T)\}$. Let f be such an edge. Clearly f is not dense. There are at most $m'(T) - 2$ colored edges that are incident to f . Hence, we can choose a free color for edge f . Now we repeat the described procedure by choosing another uncolored edge incident to e_i . Each time we choose a color that was not in the neighborhood of e_i . Hence, after coloring all the neighbors of e_i , edge e_i becomes a dominating edge. This argument applies to each e_i , $i \in \{1, \dots, m'(T)\}$, in turn.

Finally suppose that e_ℓ is uncolored for some $\ell \in \{m'(T) + 1, \dots, |E(T)|\}$. As $d(e_\ell) < m'(T) - 1$, not all of colors $1, \dots, m - 1$ appear on neighbors of e_ℓ . Hence there is some color available for e_ℓ . It follows that the constructed coloring is a b-edge coloring of T and $\varphi'(T) = m'(T) - 1$. \square

In order to deal with trees that are not pivoted we give the following definition which is closely related to pivoted trees.

Definition 2.3 *Let T be a tree and let E' be the set of dense edges of T . Suppose that E'' is a subset of E' of cardinality $m'(T)$. Then E'' encircles some edge $e \in E \setminus E''$ if:*

1. *There exist two dense edges in E'' that are incident with edge e and are not incident with each other.*
2. *Each dense edge in E'' is incident to e or to a dense edge in E'' incident to e .*
3. *Any dense edge in E'' incident to e and to another dense edge in E'' which is not incident to edge e has degree $m'(T) - 1$.*

We refer to e as an encircled edge with respect to E'' (for an example see Figure 1).




Figure 1: A pivoted tree T with $m'(T) = 6$, encircled edge e , and $\varphi'(T) = 5$

Clearly the pivot is encircled by the set of dense edges in a pivoted tree, but encircled edge of some tree T is not necessarily the pivot of T . For this observe a tree T with the encircled edge but more than $m'(T)$ dense edges. We give the additional definition that incorporates the concept of encirclement.

Definition 2.4 Let T be a tree and let E' be the set of dense edges of T . Suppose that E'' is a subset of E' of cardinality $m'(T)$. Then E'' is a good set with respect to T if:

1. Set E'' does not encircle any edge in $E \setminus E''$.
2. Any edge $e \notin E''$ with $d(e) \geq m'(T)$ is incident to some $f \in E''$ with $d(f) = m'(T) - 1$.

Lemma 2.5 Let T be a tree. If T is not a pivoted tree, then there exists a good set for T .

Proof. Let E' be the set of dense edges of T . By the definition of $m'(T)$, we may choose $E'' \subseteq E'$ with $|E''| = m'(T)$ such that every edge in $E \setminus E''$ has degree less than $m'(T)$. Let $E = \{e_1, \dots, e_{|E(T)|}\}$ be ordered in such a way that $E'' = \{e_1, \dots, e_{m'(T)}\}$.

If E'' does not encircle an edge, we are done since E'' satisfy Properties 1 and 2 of Definition 2.4. So we may assume that E'' encircles some edge $e \notin E''$. Let e_1, \dots, e_p be edges of E'' incident to e , $p \leq m'(T)$, and e_1, \dots, e_q , $q \leq p$, edges of E'' incident to e each having at least one other member of E'' , which is not incident to e , as a neighbor. Clearly $p \geq 2$ by Property 1 of Definition 2.3. Also $q \geq 1$, otherwise $p = m'(T)$ by Property 2 of Definition 2.3, and hence $d(e) \geq m'(T)$, which contradicts the choice of E'' . Thus there exists an edge e_r , $p + 1 \leq r \leq m'(T)$, incident to e_1 but not to e . Let e_i , $2 \leq i \leq p$, be an edge incident to e but not to e_1 (such an edge exists according to Property 1 of Definition 2.3).

Case 1: Suppose that edge e is dense. By the choice of E'' we have $d(e) = m'(T) - 1$. Let $W = (E'' \setminus \{e_i\}) \cup \{e\}$. Also by the choice of E'' , the only edge not in W that can have degree at least $m'(T)$ is e_i . But e_i is incident to $e \in W$, and $d(e) = m'(T) - 1$. So W satisfies Property 2 of Definition 2.4. Let $f \in E \setminus W$ be an edge different from e_i which is incident to some edge of W . Property 1 of Definition 2.3 is not fulfilled for f by W and f is not encircled by W . Also e_i is not encircled by Property 2 of Definition 2.3, since e_r is too far away from e_i . All other edges from $E \setminus W$ are not incident with any edge from W and can not be encircled by W . Hence Property 1 of Definition 2.3 is also fulfilled and W is a good set.

Case 2: Now suppose that e is not dense. If $|E'| = m'(T)$, then tree T is pivoted for e which is a contradiction. Hence $|E'| > m'(T)$ and there exists a dense edge $f \in E \setminus E''$. Let $W = (E'' \setminus \{e_1\}) \cup \{f\}$. Suppose that W encircles some edge g . At most one edge not in W lies on the path between two arbitrary non-incident edges of W , namely g . But edges $e_1 \notin W$ and $e \notin W$ lie on the path between edges $e_i \in W$ and $e_r \in W$. This contradiction implies that W satisfies Property 1 of Definition 2.4. Also, W satisfies Property 2 of Definition 2.4, since $d(e_1) = m'(T) - 1$ by Property 3 of Definition 2.3, and therefore every dense edge outside W has degree less than $m'(T)$. \square

We establish next the b-chromatic index of trees that are not pivoted.

Theorem 2.6 *If T is a tree that is not pivoted, then $\varphi(T) = m'(T)$.*

Proof. By Lemma 2.5, we may suppose that $W = \{e_1, \dots, e_{m'(T)}\}$ is a good set of $m'(T)$ dense edges of T . Such a choice is possible, since T is not pivoted. Attach color i to edge e_i , $i \in \{1, \dots, m'(T)\}$. We will show that this partial coloring can be extended to a partial b-edge coloring of T with $m'(T)$ colors, in such a way that each e_i is a dominating edge, and then to a b-edge coloring of T with the same number of colors.

Let $U = N(W) \setminus W$. We partition the set U into two subsets as follows: an edge $e \in U$ is called *inner* if there exist two edges $e_i, e_j \in W$, at distance at most 2 from each other, with e on the path between them; $e \in U$ is called *outer* otherwise. We first extend the coloring to all inner edges, then to outer edges and finally to all the remaining edges.

Suppose without loss of generality that $e_1, \dots, e_{m'(T)}$ are numbered in such a way that, if $i < j$, then e_i is incident to at least as many inner edges as e_j . Let e_1, \dots, e_p have at least two inner neighbors, e_{p+1}, \dots, e_q have one inner neighbor, and $e_{q+1}, \dots, e_{m'(T)}$ have no inner neighbors. Note that $P = \{e_1, \dots, e_p\}$ or $Q = \{e_{p+1}, \dots, e_q\}$ may be empty.

We begin by coloring uncolored inner neighbors of e_i for each i from 1 to q . Firstly, we deal with inner neighbors of e_1, \dots, e_p (assuming that $P \neq \emptyset$). For the induction step, suppose that $i \leq p$, and that inner neighbors of e_1, \dots, e_{i-1} have been colored in such a way that:

- (a) if an inner edge e is assigned its current color, say color k , during the coloring of the inner neighbors of e_j , then the path from e to e_k passes through at least one endvertex of e_j ,
- (b) no two neighbors of any edge e_j have the same color,
- (c) no two incident edges have the same color.

Note that basis of induction for e_1 follows by nothing. We show how to color uncolored inner neighbors of e_i so that these three properties continue to hold.

Let edges x_1, \dots, x_s be inner neighbors of e_i . For each $j \in \{1, \dots, s\}$, because x_j is inner, there exists an edge $e_{c_j} \in W$, $c_j \neq i$, e_{c_j} is at distance at most 1 from x_j , and x_j is on the path from e_i to e_{c_j} . Furthermore, edges e_{c_1}, \dots, e_{c_s} are distinct since there are no cycles in a tree.

Case 1: Suppose that $s > 1$. Note that c_i is the color of e_{c_i} . Let d_1, \dots, d_s be s different colors with $\{d_1, \dots, d_s\} = \{c_1, \dots, c_s\}$ and $d_i \neq c_i$ for all i . Apply color d_j to x_j , $j \in \{1, \dots, s\}$. It is straightforward to verify that Properties (a), (b), and (c) hold.

Case 2: Let $s = 1$. Because e_i has at least two inner neighbors, there is some neighbor f , which is already colored, say by the color d . Attach d to x_1 and c_1 to f . Again, it is not difficult to verify that (a), (b), and (c) hold.

Now we deal with uncolored inner neighbors of e_{p+1}, \dots, e_q (assuming that $Q \neq \emptyset$). Let z_1, \dots, z_k be those inner neighbors. Clearly $e_i, i \in \{p+1, \dots, q\}$, has at most one neighbor z_j . Thus assigning colors to z_1, \dots, z_k , at most one neighbor of each e_i is colored. It therefore suffices to ensure that, in assigning a color to each z_j , the following holds:

- (d) the partial coloring remains proper,
- (e) no e_i of degree exactly $m'(T) - 1$ has two neighbors of the same color.

Let \mathcal{C} be the set of colors defined as follows:

$$c \in \mathcal{C} \Leftrightarrow (e_c \in N(z_j)) \vee (\exists d : e_c \in N(e_d) \wedge e_d \in N(z_j) \wedge d(e_d) = m'(T) - 1),$$

where $c, d \in \{1, \dots, m'(T)\}$. If we choose a color $c \notin \mathcal{C}$, then (d) and (e) continue to hold. If $\mathcal{C} = \{1, \dots, m'(T)\}$, then z_j is encircled by W , which is a contradiction since W is a good set. Hence there is always a choice of color for z_j . Note that each $e_r, r \in \{p+1, \dots, q\}$, has only one inner neighbor colored in this way. Thus if $d(e_r) \geq m'(T)$, then at most two edges incident to e_r have the same color. All inner edges are now colored.

Next we deal with outer edges of U . Outer neighbors of $e_i, i \in \{1, \dots, m'(T)\}$ can be colored independently. Let f be an outer neighbor. The only colored edges incident to f are some dense edges and some inner edges. But they are all incident in the same endvertex of edge f , which means that every outer neighbor has at most $\Delta(T) - 1$ colored neighbors. Hence, we can choose a free color for edge f since $m'(T) > \Delta(T) - 1$. Now we repeat the described procedure by choosing another uncolored edge incident to e_i . Each time we choose a color that is not in the neighborhood of e_i . Hence, after coloring all neighbors of e_i , edge e_i becomes a dominating edge. This argument applies to each $e_i, i \in \{1, \dots, m'(T)\}$ in turn.

We may extend this partial b-edge coloring with $m'(T)$ colors to a b-edge coloring of T with $m'(T)$ colors as follows. Any remaining uncolored edge e must satisfy $d(e) \leq m'(T) - 1$. For, if $d(e) \geq m'(T)$, then by Property 2 of Definition 2.4, e is incident to some edge $f \in W$, where $d(f) = m'(T) - 1$, so that e was already colored. An edge e of degree less than $m'(T)$ cannot have neighbors colored with all colors $1, \dots, m'(T)$. Hence, there is some color available for edge e . This completes the construction of a b-edge chromatic coloring of T with $m'(T)$ colors. \square

Testing whether a tree is pivoted may be carried out in polynomial time (one needs to check for dense edges the neighborhood and the second neighborhood of each edge that is not dense). From Theorems 2.2 and 2.6 it follows that we can compute the b-chromatic index of a tree in polynomial time.

Corollary 2.7 *If T is a tree and $\mathcal{L}(T)$ is its line graph, then $\varphi(\mathcal{L}(T))$ can be determined in polynomial time.*

As we will see next, trees are already enough to show that the lower bound and the upper bound are tight with respect to the maximum degree. For this let k be a fixed positive integer and let T be a tree on n vertices with $\Delta(T) \leq k$. Clearly T has $n - 1$ edges. We construct tree T' from T as follows. Attach $k - \deg(u)$ pendant vertices to every vertex u of T . Hence T' has exactly n vertices of degree k and all other vertices of T' have degree 1. Moreover all vertices of degree k in T' form a connected subtree of T' which is T . All edges of T' that are also in T have degree $2k - 1$ and we have $n - 1$ such edges. Moreover, T' is clearly not a pivoted tree. If $n \leq \Delta(T') = k$, we have $m'(T') = \Delta(T')$ and with this also $\varphi'(T') = \Delta(T')$ by Theorem 2.6. For $\Delta(T') < n \leq 2\Delta(T')$, we have $n - 1$ edges in T' of degree $2\Delta(T') - 2$, while all other edges have degree $k - 1$. Thus $m'(T') = n - 1$ and so $\varphi'(T') = n - 1$ by Theorem 2.6. We have constructed a tree T' with $\varphi'(T') = n - 1$ for an arbitrary n with $\Delta(T') \leq n - 1 \leq 2\Delta(T') - 1$. We have proved the following theorem.

Theorem 2.8 *There exists a tree T' with maximum degree $\Delta(T')$, such that $\varphi'(T') = \ell$ for every integer ℓ and $\Delta(T') \leq \ell \leq 2\Delta(T') - 1$.*

3 Graphs with $\varphi'(G) = m'(G)$

As in the case of the b-chromatic number regular graphs play an important role also for the b-chromatic index. First reason for this is that $m'(G) = 2\Delta(G) - 1$ for any regular graph. Another reason is that we can always end the coloring by a greedy algorithm once we have a partial coloring of G in which all $2\Delta(G) - 1$ colors are realized. Thus we have $\varphi'(G) = 2\Delta(G) - 1$ for a regular graph G , if we can find such a partial coloring of G .

First recall that a graph G is of *class 1* if $\chi'(G) = \Delta(G)$ and of *class 2* if $\chi'(G) = \Delta(G) + 1$. For a vertex v of G , let $S_2(v)$ be the set of all vertices of G that are at distance 2 to v . We define the graph $G[v]$ as the subgraph of G induced by $N(v) \cup S_2(v)$.

Theorem 3.1 *Let G be a d -regular graph with $\text{diam}(G) \geq 4$ and let u and v be two vertices at distance at least 4. If $G[u]$ and $G[v]$ are class 1 graphs with $\Delta(G[u]) = \Delta(G[v]) = d - 1$, then*

$$\varphi'(G) = 2d - 1.$$

Proof. We split colors into two sets $A = \{1, \dots, d\}$ and $B = \{d+1, \dots, 2d-1\}$. Let $N(u) = \{u_1, \dots, u_d\}$ and $N(v) = \{v_1, \dots, v_d\}$. We define edge coloring $c : E(G) \rightarrow \{1, \dots, 2d - 1\}$ as follows. Let $c(u_i u) = i$ for $i \in \{1, \dots, d\}$ and $c(v_i v) = d + i$ for

$i \in \{1, \dots, d-1\}$. In addition let $c(v_d v) = 1$. Next we color all edges of $G[u]$ and $G[v]$. Since they are both class 1 graphs, there exists an edge coloring of each with $\Delta(G[u]) = \Delta(G[v]) = d-1$ colors. We use colors from B for $G[u]$ and colors from $A - \{1\}$ for $G[v]$. Note that every $u_i, i \in \{1, \dots, d\}$, has degree $d-1$ in $G[u]$ and thus have all colors from B on its incident edges. But then color i is realized on uu_i for every $i \in \{1, \dots, d\}$. Similarly every $v_i, i \in \{1, \dots, d\}$, has degree $d-1$ in $G[v]$ and thus have all colors from $A - \{1\}$ on its incident edges. But then color $j, j \in \{d+i, \dots, 2d-1\}$, is realized on vv_i for every $i \in \{1, \dots, d\}$. Hence all colors are realized and this partial coloring is a proper coloring since $d(u, v) \geq 4$. Since all colors are realized and we have $2d-1$ colors we can end the coloring by the greedy coloring. Hence c' is a b-edge coloring and $\varphi'(G) = 2d-1$. \square

Note that condition $\Delta(G[v]) = d-1$ implies that each vertex from $S_2(v)$ has a neighbor at distance 3 from v . Also for every bipartite graph G and any vertex v of G , the graph $G[v]$ is bipartite (bipartition is induced with $N(v)$ and $S_2(v)$) with $\Delta(G[v]) = d-1$. Since every bipartite graph is class 1 graph by König's Theorem, we have the next corollary.

Corollary 3.2 *If G is a bipartite d -regular graph with $\text{diam}(G) \geq 4$, then $\varphi'(G) = 2d-1$.*

In particular, for $n \geq 4$, the hypercube $Q_n = \square_{i=1}^n K_2$ has diameter n and is a bipartite n -regular graph. Hence $\varphi'(Q_n) = 2n-1$. However $\varphi'(Q_2) = 2 < 3$ and $\varphi'(Q_3) = 4 < 5$ as we will see in the next section.

One of the most important questions about the b-chromatic number is whether the Petersen graph P is the only regular graph G of girth 5 with its b-chromatic number strictly lower than $m(G)$. We will see that this is not a problem for the b-chromatic index.

Theorem 3.3 *If G is a d -regular graph with girth $g \geq 5$, then*

$$\varphi'(G) = 2d-1.$$

Proof. Let $e = uv$ be an edge in a graph G with girth $g \geq 5$ and let $N_u(e) = \{uu_2, \dots, uu_d\}$ and $N_v(e) = \{vv_{d+1}, \dots, vv_{2d-1}\}$. We define a partial edge coloring $c: E(G) \rightarrow \{1, \dots, 2d-1\}$ in the following way:

- $c(uv) = 1$,
- $c(uu_i) = i, i \in \{2, \dots, d\}$,
- $c(vv_j) = j, j \in \{d+1, \dots, 2d-1\}$.

Vertices u_i cannot be adjacent to vertices v_j , otherwise we would get a 4-cycle contrary to the assumption. Since $N(u_i) \cap N(u_j) = \emptyset$ and $N(v_i) \cap N(v_j) = \emptyset$ for all applicable indices i and j , we can continue the coloring procedure in the following way. Assign colors $\{d+1, \dots, 2d-1\}$ to all non-colored edges incident with $u_i, i \in \{2, \dots, d\}$, and colors $\{2, \dots, d\}$ to all non-colored edges incident with $v_j, j \in \{d+1, \dots, 2d-1\}$. With this partial edge coloring all colors are realized on edges uv, uu_i , and vv_j for every $i \in \{2, \dots, d\}$ and $j \in \{d+1, \dots, 2d-1\}$. Color the rest of the graph with the greedy algorithm to get a proper b-edge coloring of G with $2d-1$ colors. \square

4 Graphs with $\varphi'(G) < m'(G)$

We have shown in the previous section that for many graphs equality $\varphi'(G) = m'(G)$ holds. One can get the impression that finding $\varphi'(G)$ is not a hard problem. However as we will see in this section, things can be different when $\varphi'(G) < m'(G)$. Indeed, once this inequality holds, it seems to be very hard to find the exact value for $\varphi'(G)$.

A graph G is an *edge regular graph* or *d-edge regular graph* if all edges of G have degree d . Clearly every d -regular graph is also a $2(d-1)$ -edge regular graph and $K_{m,n}$ is a $(m+n-2)$ -edge regular graph.

Lemma 4.1 *If G is a d -edge regular graph with minimum degree $\delta \geq 4$ and $\varphi'(G) = d+1$, then at most two edges of any 4-cycle and at most two edges of any triangle realize their color in a $(d+1)$ -b-edge coloring of G .*

Proof. Let G be a d -edge regular graph with $\varphi'(G) = d+1$ and let $c : E(G) \rightarrow \{1, \dots, d+1\}$ be a b-edge-coloring of $E(G)$. Suppose that $\{u_1, u_2, u_3, u_4\}$ form a four cycle C_4 and that u_1u_2 realizes color 1. We split colors in two sets $A = \{i : c(u_1u_3) = i\}$ and $B = \{i : c(u_2u_4) = i\}$. Hence in A and in B are colors of all edges that are incident with u_1 and u_2 , respectively. Since $\varphi'(G) = d+1$, we have $A \cap B = \{1\}$ and $A \cup B = \{1, \dots, d+1\}$. In particular $c(u_2u_3) \in B$, $c(u_4u_1) \in A$, and $c(u_2u_3) \neq c(u_4u_1)$. Let $c(u_2u_3) = 2$ and $c(u_4u_1) = 4$. Moreover, if $c(u_2u_3) = 1$, then u_2u_3 and u_4u_1 do not realize colors 2 and 4, respectively, since u_1u_2 and u_3u_4 have color 1. Therefore we can assume that $c(u_3u_4) = 3$.

It is enough to show that three (or more) consecutive edges of C_4 cannot all realize their color. Suppose first that u_2u_3 realizes color 2. Color 3 must then be in A and the set C that contains colors of all edges incident with u_3 equals $\{2\} \cup (A - \{1\})$. Thus u_3u_4 does not realize color 3, since it has two edges of color 4 in its neighborhood. Also u_4u_1 does not realize color 4, since it has two edges of color 3 in its neighborhood.

Let now $u_1u_2u_3$ be a triangle in G . Set $c(u_1u_2) = 1$, $c(u_2u_3) = 2$, and $c(u_3u_1) = 3$. Suppose that u_1u_2 and u_2u_3 realize colors 1 and 2, respectively. Let A and B be

as before. Then $C = \{i : c(u_3w) = i\}$ equals to $\{2\} \cup (A - \{1\})$ since $\varphi'(G) = d + 1$ and u_1u_2 and u_2u_3 realize their colors. Hence u_1u_3 does not realize its color since no colors from $B - \{1, 2\}$ are in the neighborhood of u_1u_3 . \square

Note, from the above proof, that we have two possibilities. Namely, realizable edges of C_4 are either two consecutive edges of C_4 or two opposite edges of C_4 . Also C_4 is not necessarily an induced cycle. But if C_4 is not induced, we need the additional condition $\delta \geq 4$ of Lemma 4.1. If $u_1u_3 \in E(G)$, then $c(u_1u_3) \neq 3$ and we need an additional edge incident with u_1 so that color 3 is in A . Hence if we demand an induced cycle C_4 , we do not need the condition $\delta \geq 4$ anymore.

By observing $\varphi'(K_n)$ for small n it seems that it equals to $\chi'(K_n)$. Namely, it is not hard to see that $\varphi'(K_3) = 3$, $\varphi'(K_4) = 3$ and $\varphi'(K_5) = 5$. However we have $\varphi'(K_6) = 6$ which breaks the above suggestion. Indeed, the following 6-b-edge-coloring of K_6 is due to Stephan Brandt (personal communication). Color consecutive edges of C_6 by 1,2,3,1,2,3. The remaining edges of $K_6 - E(C_6)$ form a triple perfect matching. Edges of each of this matching receive the same color 4,5, and 6.

Proposition 4.2 *If $n \geq 4$, then $\varphi'(K_n) < m'(K_n)$.*

Proof. As mentioned above $\varphi'(K_4) = 3 < 5 = m'(K_4)$. Now let $n \geq 5$. Suppose that $\varphi'(K_n) \geq m'(K_n) = 2n - 3$ and let c be a b-edge coloring of $E(K_n)$ with $2n - 3$ colors. Let H be a spanning subgraph of K_n whose edges are realizable edges of c . By Lemma 4.1 H has no triangle and no four cycle. Moreover, H is a forest. Namely, if C_k , $k > 4$, is in H , then three consecutive edges of C_k are all realizable on a four cycle in contradiction to Theorem 4.1. This forest has the most edges if it is a tree T . But a spanning tree T has at most $n - 1$ edges which is a contradiction. \square

For complete bipartite graphs, recall the well-known result (see [13, Proposition 1.2]) that $\mathcal{L}(K_{p,r}) = K_p \square K_r$. Kouider and Mahéo have shown in [19] that $\varphi(K_p \square K_r) = r$ whenever $r \geq p(p - 1)$ and for $p \leq r < p(p - 1)$ we have $r \leq \varphi(K_p \square K_r) \leq p(p - 1)$. Hence we have $\varphi'(K_{p,r}) = r$ for $r \geq p(p - 1)$ and $r \leq \varphi'(K_{p,r}) \leq p(p - 1)$ for $p \leq r < p(p - 1)$. In particular, for $p = 2$, we have $\varphi'(K_{2,r}) = r$ for every $r \geq 2$. We can immediately improve the upper bound in many cases since $K_{p,r}$ is an edge regular graph with $m'(K_{p,r}) = p + r - 1$. Hence

$$\varphi'(K_{p,r}) \leq \min\{p(p - 1), p + r - 1\} \text{ for } p \leq r < p(p - 1).$$

To improve this we need some further arguments.

Proposition 4.3 *If $r \geq p \geq 3$, then $\varphi'(K_{p,r}) < m'(K_{p,r})$.*

Proof. If $r \geq p(p-1)$ we are done by results from [19] (see the above discussion). Thus suppose that $\varphi'(K_{p,r}) = m'(K_{p,r}) = p+r-1$ and let c be a b-edge-coloring of $E(K_{p,r})$ with $p+r-1$ colors. Let H be a spanning subgraph of $K_{p,r}$ whose edges are realizable edges of c . Again H is a forest, since every cycle C_k , $k \geq 4$, induce a four cycle in $K_{p,r}$ with three realizable edges, which is not possible by Lemma 4.1. Moreover, if edges uv and vw are realizable for some colors, then no edge ux or wx is realizable, since it is on a common four cycle $uvwxu$. Hence we must find a cover of $K_{p,r}$ by stars with maximum number of edges and every edge of this stars can be a realizable edge. Clearly this number is $p+r-2$ if we take two stars $K_{1,r-1}$ and $K_{1,p-1}$. Thus we have $p+r-2$ candidates for realizable edges, which is a contradiction, since we need at least $p+r-1$ candidates. \square

We believe that this upper bound can be lowered by one in general, but not more as can be seen from the following schemes. On the scheme we present the b-edge colorings of $K_{3,3}$, $K_{4,4}$, and $K_{5,5}$ with 3, 5, and 7 colors, respectively:

$$K_{3,3} : \begin{array}{ccc} \tilde{1}\tilde{2}\tilde{3} & 231 & 312 \\ \tilde{1}23 & \tilde{2}31 & \tilde{3}12 \end{array} ; K_{4,4} : \begin{array}{cccc} \tilde{1}\tilde{2}\tilde{3}\tilde{4} & \tilde{4}523 & \tilde{5}342 & 2451 \\ \tilde{1}\tilde{4}\tilde{5}2 & \tilde{2}534 & \tilde{3}245 & 4321 \end{array} ;$$

$$K_{5,5} : \begin{array}{ccccc} \tilde{1}\tilde{2}\tilde{3}\tilde{4}\tilde{7} & \tilde{5}6734 & \tilde{6}7413 & 73162 & 256\tilde{7}1 \\ \tilde{1}\tilde{5}\tilde{6}\tilde{7}2 & \tilde{2}6735 & \tilde{3}7416 & \tilde{4}316\tilde{7} & 54321 \end{array} .$$

Here 12347 in first line and first column of $K_{5,5}$ represents colors of u_1w_1 , u_1w_2 , u_1w_3 , u_1w_4 , and u_1w_5 , respectively, and \tilde{k} means that this edge realizes color k . Unfortunately we could not find a pattern for every graph $K_{n,n}$. Next we show that there exists no 4-b-edge coloring of $K_{3,3}$, which is a difficult task already, since we cannot use Lemma 4.1 anymore.

Proposition 4.4 $\varphi'(K_{3,3}) = 3$.

Proof. Suppose that there exists a 4-b-edge coloring of $K_{3,3}$ and let c be a b-edge-coloring of $E(K_{3,3})$ with 4 colors. Let H be a spanning subgraph of $K_{3,3}$ whose edges are realizable edges of c . We will analyze all different possibilities for H . If H is isomorphic to a forest of two stars $K_{1,2}$, we have a contradiction, since the edge between the centers of these two stars cannot be colored by any of the four colors. If H contains C_4 as a subgraph, we have a contradiction since the edges between vertices of $K_{3,3}$ that are not in C_4 cannot be colored. If H is isomorphic to $P_4 \cup K_2$, the middle edge of P_4 is not realizable, since the color of K_2 cannot be in its neighborhood. If there exists a path P_5 in H and an isolated vertex x , we first concentrate on the first and last edge e and f , respectively, of P_5 . It is easy to see that all edges, except the edge between the middle vertex of P_5 and x , must be colored, so that e and f are realizable edges. In that case the middle edges of P_5 are not realizable, since only one edge remains and each of them needs one additional

color. Finally, suppose that H contains one component that is isomorphic to $K_{1,3}$ in which one edge is subdivided, and one isolated vertex. It is easy to see, that edges of $K_{1,3}$ that were not subdivided cannot both be realizable, since only one can have the color of the other pendant edge in its neighborhood. \square

It is straightforward to see that a k -partite complete graph K_{n_1, \dots, n_k} is an edge regular graph if and only if $n_1 = \dots = n_k = n$. Then K_{n_1, \dots, n_k} is $(k-1)n$ -regular graph and we will use notation K_n^k for this graph.

Proposition 4.5 *If $k \geq 3$ and $n \geq 2$ are positive integers then $\varphi'(K_n^k) < m'(K_n^k)$.*

Proof. Suppose that $\varphi'(K_n^k) = m'(K_n^k) = 2(k-1)n - 1$ and let c be a b -edge-coloring of $E(K_n^k)$ with $2(k-1)n - 1$ colors. Let H be a spanning subgraph of K_n^k whose edges are realizable edges of c . By Lemma 4.1 H is a forest. In contrast to the proof of Proposition 4.3, here we can have a component different than a star. Clearly we have more edges if the number of components of H is smaller. Nevertheless, even if H is a tree, we have $kn - 1$ edges in H , which is not enough, since $k \geq 3$. \square

Let G be a regular graph. In view of Theorem 3.1, one can expect more graphs with $\varphi'(G) < m'(G)$ among those with small diameter (< 4). Such are Hamming graphs with two or three factors. It seems that it is quite hard to give an exact answer for $\varphi'(K_p \square K_r)$. The impression is that, if p and r are ‘‘large’’ enough, we have equality between φ' and m' . For this note that from Lemma 4.1 it easily follows that $\varphi'(K_2 \square K_3) < m'(K_2 \square K_3) = 5$, but in the following scheme there is a 4- b -edge coloring of $K_2 \square K_3$, which yields $\varphi'(K_2 \square K_3) = 4$. On this scheme also an 11- b -edge coloring of $K_3 \square K_5$ is presented (note that $m'(K_3 \square K_5) = 11$).

$$\begin{array}{rcccl}
 & & & \tilde{1}\tilde{2}\tilde{3}\tilde{4} & 7 & 11 & 8 & 8 & 11 & 7 & - \\
 & & & 3 & 10 & 10 & 10 & 10 & 10 & & \\
 K_2 \square K_3 : & \tilde{2}\tilde{4} & \tilde{3} & - & \tilde{5}\tilde{6}\tilde{7}\tilde{8} & 3 & 4 & 11 & 11 & 4 & 3 & - \\
 & \tilde{1} & 1 & 1 & ; & 1 & 2 & 2 & 2 & 2 & 2 & \cdot \\
 & \tilde{3}\tilde{4} & 2 & - & \tilde{9}\tilde{10}\tilde{11}\tilde{13} & 4 & 8 & 7 & 7 & 8 & 4 & - \\
 & & & & 5 & 6 & 6 & 6 & 6 & 6 & &
 \end{array}$$

Again \tilde{k} is the edge that realizes color k . Every odd line represents colors of edges of a layer of the second factor (K_3 or K_5), while every even line represents colors of edges of the first factor (K_2 or K_3) that projects to the same edge. Furthermore, for $V(K_n) = \{1, \dots, n\}$, in every odd line numbers that are written in i -th column present consecutive colors of edges $(i, i+1), \dots, (i, n)$ for $i \in \{1, \dots, n-1\}$.

For $Q_3 = K_2 \square K_2 \square K_2$ it is easy to see, with the use of Lemma 4.1, that $\varphi'(Q_3) < m'(Q_3) = 5$. Also it is easy to construct a 4- b -edge coloring of Q_3 (every perfect matching receives all four colors). Hence $\varphi'(Q_3) = 4$. In general for $K_p \square K_q \square K_r$, $p \leq q \leq r$, it seems that if $q > 2$ we can expect equality between φ'

and m' . On the other hand we do not dare to predict what happens if $p = q = 2$. Namely, if φ' is always less than m' , or there exists an integer k , such that equality holds for every $r \geq k$, or equality holds for some integers and for some not.

5 3-regular graphs

We already know that if G is a d -regular graph, then $\varphi'(G) \leq 2d - 1$. Many of d -regular graphs achieve this trivial upper bound by Theorems 3.1 and 3.3. Determining $\varphi'(G)$ is equivalent to determining $\varphi(\mathcal{L}(G))$ which is also a regular graph. According to the theorem of Kratochvíl, Tuza, and Voigt [22] there are only a finite number of regular graphs with φ smaller than the trivial upper bound. Hence there are only a finite number of d -regular graphs G with $\varphi'(G) < 2d - 1$. Can we find them all? In [16] all exceptions for the b -chromatic number were described for the smallest nontrivial d -regular graphs, namely for cubic graphs. We follow their approach also for the b -chromatic index of cubic graphs. The following lemma provides a useful tool to decrease the number of cases to be treated.

Lemma 5.1 *Let G be a cubic graph. If G has an induced cycle C_5 or an induced path P_6 , then $\varphi'(G) = 5$.*

Proof. Suppose that a cubic graph G has an induced cycle $C_5 = x_1x_2x_3x_4x_5$. Color edges of C_5 by colors $1, \dots, 5$. Since C_5 is induced, there are no edges between its vertices. For any $i \in \{1, 2, 3, 4, 5\}$ denote with y_i the third vertex to which the vertex x_i is adjacent, see Figure 2. Note that some y_i s might represent the same vertex. According to Figure 2 it is obvious that all edges of C_5 are realizable. We can complete the coloring to whole G by the greedy algorithm.




Figure 2: Induced subgraph C_5

Suppose now that G has an induced path $P_6 = x_1x_2x_3x_4x_5x_6$. Color edges of P_6 by colors $1, \dots, 5$. Since P_6 is induced, there are no edges between its vertices.

Let y_0 and y_1 be two additional neighbors of vertex x_1 . Further, let y_6 and y_7 be two additional neighbors of vertex x_6 . For $i \in \{2, 3, 4, 5\}$ the vertex x_i has only one additional neighbor y_i . See Figure 3 for this notations. Note that some y_i s might represent the same vertex. Note that at most one of y_0 and y_1 can be equal to y_3 and at most one of y_6 and y_7 can be equal to y_4 . If this is the case, then incident edges receive different colors (see the lower graph on Figure 3). Now even if one of y_0 or y_1 equals to y_6 or y_7 (see the lower graph on Figure 3), we can properly color edges so that all colors on P_6 are realizable. The rest of G can again be colored by the greedy algorithm. \square




Figure 3: Induced subgraph P_6

Now we can prove the main theorem of this section.

Theorem 5.2 *Let G be a connected cubic graph. Then $\varphi'(G) = 5$ if and only if G is not isomorphic to K_4 , $K_3 \square K_2$, $K_{3,3}$, or Q_3 . Moreover, $\varphi'(K_4) = 3$, $\varphi'(K_{3,3}) = 3$, $\varphi'(K_3 \square K_2) = 4$, and $\varphi'(Q_3) = 4$.*

Proof. We have already seen that $\varphi'(K_4) = 3$, $\varphi'(K_{3,3}) = 3$, $\varphi'(K_3 \square K_2) = 4$, and $\varphi'(Q_3) = 4$.

Let now G be a cubic graph not isomorphic to K_4 , $K_3 \square K_2$, $K_{3,3}$, or Q_3 . The proof of this direction is constructive and leads to an algorithm that finds appropriate subgraphs. Moreover, according to Lemma 5.1 we wish to find an induced cycle C_5 or an induced path P_6 in analyzed cases.

We will analyze cases with respect to the girth of a given cubic graph G . Let g be its girth and let C be a g -cycle of G . For $i \geq 1$, let

$$D_i = \{v \in V(G) : d(v, C) = i\}$$

be the i -th distance level from C . According to Theorem 3.3 we only need to consider cases where $g = 3$ and $g = 4$. In the following figures we usually present only a part

of the graph in which we find an induced C_5 or P_6 , since the rest can be colored by the greedy algorithm. Also we do not draw all possibilities that have the same induced C_5 or P_6 , but just one representative.

Case 1: $g = 3$.

In this case C is a triangle. We distinguish subcases with respect to the size of D_1 .

Case 1.1: $|D_1| = 1$.

In this subcase $G \cong K_4$ which is forbidden.

Case 1.2: $|D_1| = 2$.

Let $D_1 = \{x, y\}$. Note that there is a unique way (up to the isomorphism) how x and y are adjacent with the vertices of C . Assume without loss of generality that x has two neighbors in C (and hence y has one).

Suppose first that $xy \notin E(G)$. If x and y have a common neighbor in D_2 , then we have an induced C_5 , see the left graph on Figure 4. If x and y have no common neighbor in D_2 , then $|D_2| = 3$. If vertices of D_2 induce a triangle, then there is no vertex in D_3 and this graph has no induced C_5 or P_6 , but there is a 5-b-edge coloring on the second graph of Figure 4. If D_2 does not induce a triangle, then $|D_3| \geq 1$ and one neighbor of y has a neighbor in D_3 . This yields an induced P_6 , see the third graph of Figure 4. Finally if $xy \in E(G)$, then $|D_2| = 1$, $|D_3| = 2$, and $|D_4| \geq 1$. In all cases there exists an edge from D_3 to D_4 as denoted on the right graph of Figure 4, and therefore we have an induced P_6 .




Figure 4: Subgraphs with $g = 3$ and $|D_1| = 2$

Case 1.3: $|D_1| = 3$.

In this subcase each vertex from C has its own private neighbor in D_1 . If vertices of D_1 induced a triangle, we get the forbidden graph $K_3 \square K_2$. Hence suppose that D_1 does not induce a triangle and we always have two nonadjacent vertices in D_1 . If two nonadjacent vertices of D_1 have a common neighbor in D_2 , then there exists an induced C_5 , see the left graph of Figure 5. If D_1 induces $3K_1$ and no two vertices of D_1 have a common neighbor in D_2 , then we have $|D_2| = 6$, and there exists two nonadjacent vertices in D_2 with different neighbors in D_1 . They form an induced P_6 , see the second graph of Figure 5 for one possibility. Suppose now that D_1 induces $K_2 \cup K_1$ and denote K_1 by x . If a neighbor of x in D_2 coincides with a neighbor of some other vertex of D_1 , we get an induced C_5 , see the third graph of Figure 5. If a neighbor of x in D_2 is nonadjacent with a vertex of D_2 , which is nonadjacent to x , we have an induced P_6 , see the fourth graph of Figure 5 (dotted line means there

is no edge). Otherwise both neighbors of x in D_2 are adjacent to both neighbors in D_2 of K_2 in D_1 . This graph contains an induced C_5 , see the fifth graph of Figure 5. Finally, if D_1 induces P_3 , we have an induced C_5 , see the last graph of Figure 5.




Figure 5: Subgraphs with $g = 3$ and $|D_1| = 3$

Case 2: $g = 4$.

Now C is a square. Note that two adjacent vertices of C have no common neighbor in D_1 . Once more we distinguish subcases with respect to the size of D_1 . It is obvious that the case $|D_1| = 1$ is not possible.

Case 2.1: $|D_1| = 2$.

If vertices of D_1 are adjacent, we obtain $K_{3,3}$ which is forbidden. We have an induced C_5 if vertices of D_1 have a common neighbor in D_2 , see the left graph of Figure 6. If vertices of D_1 have no common neighbor in D_2 , we have $|D_2| = 2$. If these two vertices are not adjacent, we have an induced P_6 , see the middle graph of Figure 6. Otherwise, they are adjacent and $|D_3| \geq 1$, which again results in an induced P_6 , see the right graph of Figure 6.




Figure 6: Subgraphs with $g = 4$ and $|D_1| = 2$

Case 2.2: $|D_1| = 3$.

Here two nonadjacent vertices of C must have a common neighbor x in D_1 , while the other two have exactly one neighbor y and w in D_1 . If $yw \in E(G)$, we have an induced C_5 , see the first graph of Figure 7. If $xy \in E(G)$ (and analogue when $xw \in E(G)$), we have $|D_2| \geq 2$. This yields an induced P_6 , since w has two neighbors in D_2 and only one of them can be equal to the neighbor of y in D_2 , see the second graph of Figure 7. Hence no two vertices of D_1 are adjacent. If x and y have a common neighbor in D_2 , then these vertices form an induced C_5 , see the third graph of Figure 7. If a neighbor of x in D_2 is nonadjacent to some other vertex of D_2 , we have an induced P_6 , see the fourth graph of Figure 7. Otherwise we have the last graph of Figure 7 which also contains an induced P_6 .

Case 2.3: $|D_1| = 4$.

Every vertex of C has his own private neighbor in D_1 . Denote them by x_1, x_2, x_3 , and x_4 . If $x_1x_3 \in E(G)$ (or analogue $x_2x_4 \in E(G)$), we have an induced C_5 , see the




Figure 7: Subgraphs with $g = 4$ and $|D_1| = 3$

left graph of Figure 8. Now D_1 does not induce a cycle C_4 , since we have either the previous possibility or Q_3 which is forbidden. If D_1 induces paths P_2 , P_3 , $2P_2$, or P_4 , see the second, third, fourth, and fifth graph, respectively, of the Figure 8 for an induced P_6 , which starts and ends in nonadjacent vertices of D_1 . Thus it remains to deal with the cases when there are no edges between vertices of D_1 . If x_i and $x_{(i+1)\bmod 4}$ have a common neighbor, for some $i \in \{1, 2, 3, 4\}$, we have an induced C_5 , see the right lower graph of Figure 8 for one possibility. If x_i and $x_{(i+2)\bmod 4}$ do not have a common neighbor, for some $i \in \{1, 2, 3, 4\}$, we have an induced P_6 , see the left lower graph of Figure 8 for one possibility. Otherwise x_1 and x_3 have a common neighbor in D_2 . Note that x_1 and x_2 , x_2 and x_3 have no common neighbor in D_2 . Again we have an induced P_6 , see the middle lower graph of Figure 8.




Figure 8: Subgraphs with $g = 4$ and $|D_1| = 4$

□

Acknowledgements

This work has been financed by ARRS Slovenia under the grant P1-0297 and within the EUROCORES Programme EUROGIGA (project GReGAS) of the European Science Foundation.

References

- [1] R. Balakrishnan, S. Francis Raj, Bounds for the b -chromatic number of the Mycielskian of some families of graphs, to appear in Ars Combin.

- [2] R. Balakrishnan, S. Francis Raj, Bounds for the b -chromatic number of $G - v$, Discrete Appl. Math. (2011) Doi:10.1016/j.dam.2011.08.022.
- [3] R. Balakrishnan, S. Francis Raj, T. Kavaskar, Coloring the Mycielskian, Proc. Int. Conf. ICDM (2008) 53–57.
- [4] D. Barth, J. Cohen, T. Faik, On the b -continuity property of graphs, Discrete Appl. Math. 155 (2007) 1761–1768.
- [5] S. Cabello, M. Jakovac, On the b -chromatic number of regular graphs, Discrete Appl. Math. 159 (2011) 1303–1310.
- [6] F. Chaouche, A. Berrachedi, Some bounds for the b -chromatic number of a generalized Hamming graphs, Far East J. Appl. Math. 26 (2007) 375–391.
- [7] S. Corteel, M. Valencia-Pabon, J-C. Vera, On approximating the b -chromatic number, Discrete Appl. Math. 146 (2005) 106–110.
- [8] B. Effantin, The b -chromatic number of power graphs of complete caterpillars, J. Discrete Math. Sci. Cryptogr. 8 (2005) 483–502.
- [9] B. Effantin, H. Kheddouci, The b -chromatic number of some power graphs, Discrete Math. Theor. Comput. Sci. 6 (2003) 45–54.
- [10] B. Effantin, H. Kheddouci, Exact values for the b -chromatic number of a power complete k -ary tree, J. Discrete Math. Sci. Cryptogr. 8 (2005) 117–129.
- [11] R. Hammack, W. Imrich, S. Klavžar, Handbook of Product Graphs, Second Edition, Discrete Mathematics and Its Applications, CRC Press, Boca Raton, 2011.
- [12] C.T. Hoang, M. Kouider, On the b -dominating coloring of graphs, Discrete Appl. Math. 152 (2005) 176–186.
- [13] W. Imrich, S. Klavžar, D.F. Rall, Topics in Graph Theory: Graphs and Their Cartesian Product, A.K. Peters, Wellesley, MA, 2008.
- [14] A. Iranmanesh, I. Gutman, O. Khormali, A. Mahmiani, The edge version of the Wiener index, MATCH Commun. Math. Comput. Chem. 61 (2009) 663–672.
- [15] R.W. Irving, D.F. Manlove, The b -chromatic number of a graph, Discrete Appl. Math. 91 (1999) 127–141.
- [16] M. Jakovac, S. Klavžar, The b -chromatic number of cubic graphs, Graphs Combin. 26 (2010) 107–118.
- [17] M. Jakovac, I. Peterin, On the b -chromatic number of some products, Studia Sci. Math. Hungar. 49 (2012) 156–169.

- [18] M. Kouider, *b*-chromatic number of a graph, subgraphs and degrees, Res. Rep. 1392, LRI, Univ. Orsay, France, 2004.
- [19] M. Kouider, M. Mahéo, Some bounds for the *b*-chromatic number of a graph, Discrete Math. 256 (2002) 267–277.
- [20] M. Kouider, M. Mahéo, The *b*-chromatic number of the Cartesian product of two graphs, Studia Sci. Math. Hungar. 44 (2007) 49–55.
- [21] M. Kouider, M. Zaker, Bounds for the *b*-chromatic number of some families of graphs, Discrete Math. 306 (2006) 617–623.
- [22] J. Kratochvíl, Z. Tuza, M. Voigt, On the *b*-chromatic number of graphs, Lecture Notes in Comput. Sci. 2573 (2002) 310–320.
- [23] C. V. G. C. Lima, N. A. Martins, L. Sampaio, M. C. Santos, A. Silva, *b*-chromatic index of graphs, manuscript 2013.